

UDC 599.4(611)

PRELIMINARY STUDY OF HABITAT-RELATED BAT FAUNA OF MASTOUTA-BISHSHOUK REGION (NORTHWEST TUNISIA)

R. Dalhoumi¹, A. Nefla², W. Bedoui³, R. Ouni⁴, P. Aissa¹, S. Aulagnier⁵

¹LR01ES14 Bio Surveillance de l'Environnement, Faculté des sciences de Bizerte,
Université de Carthage, 7021 Zarzouna (Tunisia)

E-mail: ridhadalhoumi@gmail.com

²Département de Biologie, Faculté des Sciences de Tunis,
Université de Tunis, El Manar II 2092 Tunis (Tunisia)

³Laboratoire de Parasitologie, Ecole Nationale de Médecine Vétérinaire,
Université de la Manouba. 2020 Sidi Thabet, Tunis (Tunisia)

⁴Sidi Thabet, Bp 23, 2020, Tunis (Tunisia)

⁵Comportement et Écologie de la Faune Sauvage, I.N.R.A., Université de Toulouse, CS 52627,
31326 Castanet Tolosan cedex (France)

E-mail: Stephane.Aulagnier@toulouse.inra.fr

Preliminary Study of Habitat-Related Bat Fauna of Mastouta-Bishshouk Region (Northwest Tunisia).

Dalhoumi, R., Nefla, A., Bedoui, W., Ouni, R., Aissa, P., Aulagnier, S. — In order to define appropriate conservation measures in northern Tunisia we surveyed the bat fauna of Mastouta-Bishshouk region in both roosting sites and foraging areas. A total of 11 species was recorded. We found only three occupied roosts including a maternity colony of *Rhinolophus ferrumequinum*, *Miniopterus schreibersii* and *Myotis cappacinii* in a train abandoned tunnel. A tunnel of water channel hosted *Rhinolophus mehelyi*, *M. schreibersii* and *M. punicus*. Echolocation calls were recorded at a bridge over Beja Wadi, at an artificial basin, and in crop fields. Bat activity was the highest at the bridge in late March, and null in the harvested crop fields in late August. *Pipistrellus kuhlii* was the most active bat species in the three sites, *Pipistrellus pipistrellus* was the second most active species, followed by *Eptesicus isabellinus*. *Plecotus gaisleri* and *Myotis punicus* were detected only once at the bridge and the basin respectively. This bat assemblage is representative of the bat fauna of northern Tunisia. Roost and foraging area surveys strongly benefit acoustic recording, even in winter when some bats remain active. The rare local underground roosts should be protected from human disturbance and the water quality of Majerda Wadi should be improved.

Key words. Chiroptera, water, tunnel, Mediterranean, North Africa.

Introduction

Terrestrial organisms live in an environment that is subject to profound rhythmic alteration including a daily alternation of light and darkness that affects their circadian activity (Erkert, 1982). As strictly nocturnal animals, bats need a roost where they can rest during the day, fulfilling their physiological demands and safe from predators (Findley, 1993; Neuweiler, 2000). They also need foraging areas related to their dietary diversity to be visited during the night (Altringham, 2011). Then roosting habits and feeding ecology require different complementary habitats, influencing local and global distribution of bats (Altringham, 2011). So conservation efforts should target both roosting and foraging habitats. Before the use of ultrasound detection that enabled field identification of foraging bats, most studies relied on identifying and counting bats in their roost in Europe (Racey, 1998), and in all other parts of the World.

In Tunisia, since the first data published by Hartmann (1868), the bat fauna has been sporadically investigated (see Dalhoumi et al., 2011). In northern Tunisia, most data come from direct observations in a small number of roost sites (Blanc, 1935; Cockrum, 1976; Deleuil & Labbe, 1955; Gadeau de Kerville, 1908; Noblet & Nefzi, 1991). Among the most recent studies in northern Tunisia, Zava & Massetti (2007) listed eight bat species in El Feidja National Park during an expedition in late May 2000 and Puechmaille et al. (2012) reported 14 bat species in the northern part of Tunisia as the result of expeditions between May 2008 and July 2012. None of these surveys provide information on habitat use by bats. The first data from acoustic sampling were reported by Rebelo & Brito (2006) who focused on bat activity in Saharan areas. Then, Puechmaille et al. (2012) used acoustic sampling to survey Tunisian bat fauna. More recently, and for the first time, Dalhoumi et al. (2014, 2016, 2017 a, b) studied the seasonal patterns of bat activity and habitat use in central Tunisia. However the later studies have been conducted in arid zones whereas northern Tunisia is mainly an agricultural land occupied by crop fields. Natural vegetation is confined to high hills and mountain regions. During the last decade and particularly after the political turmoil of 2011, natural habitats suffered serious degradations caused by intensification of land use, extensive stockbreeding, cutting and burning of forests that destroyed 2000 ha in 2012 (Chriha & Sghari, 2013). These damages resulted in fragmenting natural habitats which may be detrimental to certain bat species using linear features connecting foraging areas (Russ & Montgomery 2002). They might have altered the trophic organization of communities (Hill et al., 2011), affected the activity level of many bat species (Bright, 1993), and even led to a decline of other ones (Castro & Michalski, 2014; Wickramasinghe et al., 2003).




Fig. 1. Bat sampling sites in the Mastouta-Bishshouk region (Beja, Northwest Tunisia). 1 — Bridge of Beja Wadi, 2 — Artificial basin, 3 — Crop fields, 4 — Tunnel of water channel, 5 — Train abandoned tunnel.

In order to define appropriate conservation measures for bats in northern Tunisia, we surveyed the area along the Majerda Wadi to locate roosts, and investigated for the first time three sites corresponding to three foraging habitat types.

Study area

The study was conducted in Mastouata-Bishshouk region (Beja governorate, Tunisia) (fig. 1) on the western part of the freshwater lake of Sidi Salem Dam that was built in 1981 across the Majerda Wadi. The main habitat types are wetlands crossed from east to west by Majerda Wadi surrounded by a mosaic of crop fields and human settlements in the plain, Mediterranean shrublands and woodlands on top of hills up to 300–350 m high; the area also includes cliffs, rocky habitats, water springs and wadis. The natural vegetation is restricted to wetlands with *Phragmites communis*, *Tamarix africana*, *Typha latifolia*, *Scirpus maritimus* and *Jonca maritimus*, river banks with *Acacia cyclops*, and top of the hills with *Rosmarinus officinalis*, *Calycotoma vilosa*, *Thymus capitatus*, *Pinus* spp. and *Cupressus sempervirens*. The plains and low hills are occupied by cereal fields. The climate is semi-arid with monthly mean temperature ranging from 10 to 30 °C, and an average annual rainfall of 500 mm.

Material and methods

Bats were surveyed in March 2016, August 2016 and February 2017 in order to sample both winter and summer assemblages by roost search and foraging habitat investigation.

Roost survey

We visited two tunnels and various caves in the study area. The tunnel of water channels (about 1 km long) was inspected only in late February 2017 when its northwestern entrance was closed by straw bales. Some specimens were captured and identified using Dietz et al. (2007) and Aulagnier et al. (2009) and immediately released. The train abandoned tunnel (about 300 m in length) was inspected only in late August 2016 when the soil was dry following the unusual low level of Majerda Wadi during this summer. Echolocation calls of roosting bats were recorded during a night and a diurnal visits in August 2016, at their emergence from the northern entrance. In February 2017, they were recorded using a SM2+ bat detector (Wildlife Acoustics, Inc. Concord, Massachusetts, U.S.A.) and identified as previously described.

Foraging habitat investigation

At each period, we recorded bat activity at one fixed point near the bridge of Baja Wadi, at the artificial basin near the strait of El Hajama, and along one transect of 300–400 m in crop fields near Dahmania (fig 1). Each of the nine sessions, one per period and site, started 15 min after sunset and lasted for 2 hours. Bat calls were detected and recorded with a D240x ultrasound detector (Pettersson Elektronik AB, Uppsala, Sweden) connected to a portable digital tape (Edirol R-09HR; Roland Corporation, Shanghai, China). Calls were subsequently analyzed using the software BatSound, v.3.10 (Petterson Electronik AB). We used a simple frequency of 44.100 samples/s with 16 bits/sample and selected a 512-pt FFT with a Hamming window for analysis (Russ, 1999). We counted one pass every 37.4 s (3.4 s: recording time of the bat detector; 34 s: recording time of the portable digital tape) for each individual. In February 2017 the SM2+ bat detector was also used at the artificial basin together with the active survey with the D240x.

Recorded calls were identified to species level using shape and several call parameters (Barataud, 2012). Echolocation calls of *Tadarida teniotis*, *Miniopterus schreibersii*, *Eptesicus isabellinus*, *Plecotus gaisleri*, *Pipistrellus kuhlii*, *Myotis punicus* and *Myotis capaccinii* were identified using reference calls collected in different places of Tunisia (mainly near Mastouata, and in Jbel Erwa, Bou Hedma, Jbel Mghilla and Dghoumous National Parks). According to Russo and Jones (2002), *Pipistrellus pipistrellus* was identified by their shallow FM/QCF calls at frequency of maximum energy of 46.9 ± 1.8 kHz. *Rhinolophus ferrumequinum* and *Rhinolophus hipposideros* were easily identified as they produced calls with a long constant frequency part differing by peak frequency, between 85 and 88 kHz for *R. ferrumequinum*, over 112 kHz for *R. hipposideros* (Puechmaille et al., 2012).

Activity (number of passes) among the three sites was compared within each session by a chi-square test.

Results

A total of 11 species was recorded in the study area, 3 in the tunnel of water channel, 5 in the train abandoned tunnel, whereas 4 were recorded at the bridge of Beja Wadi, 6 at the artificial basin, and 2 over the crop fields (table 1). Specific richness and bat activity varied according to site and season.

Roosting bats

Among the five subterranean sites reported in the study area, we found guano in the train abandoned tunnel (fig. 1), in the tunnel of water channel and in a crack on Yagouma's Island where no bat was observed in February 2017. A total of nine species were recorded. In the train abandoned tunnel, we detected echolocation calls of *Tadarida teniotis*, *Miniopterus schreibersii*, *E. isabellinus*, *P. kuhlii*, and *M. capaccinii* during a night pass and calls of *Rhinolophus ferrumequinum*, *M. schreibersii* and *M. capaccinii* during a diurnal pass in August 2016. We then found a maternity colony of *R. ferrumequinum*, *M. schreibersii* and *M. capaccinii* in several small cracks of this tunnel's ceiling. In February 2017 we captured *Rhinolophus mehelyi*, *M. schreibersii* and *M. punicus* in the tunnel of water channel, and the SM2+ bat detector recorded calls of *M. capaccinii*, *T. teniotis*, *M. schreibersii*, *E. isabellinus*, *P. kuhlii* and *R. hipposideros* at the northwestern entrance of the train abandoned tunnel.

Foraging bats

A total of 1073 echolocation calls were recorded during the whole sessions (table 1), and seven bat species were recorded. *P. kuhlii* was the most frequent and active species. It wasn't detected only in harvested fields of Hanshir Bishshouk (August 2016). *P. pipistrellus* was the second most active species and was reported particularly at the bridge of Baja Wadi together with *E. isabellinus*. Other species were rarely detected. Signals of *R. hipposideros*, *M. capaccinii* and *M. punicus* were recorded only near the artificial basin, and those of *P. gaisleri* only at the bridge (table 1).

Table 1. Acoustic activity of bats recorded at three sites of the Mastouta-Bishshouk region (Beja, Northwest Tunisia)

Date	March 2016			August 2016			February 2017			Total
Locality	Bridge	Basin	Fields	Bridge	Basin	Fields	Bridge	Basin	Fields	
<i>R. ferrumequinum</i>	-	-	-	-	-	-	-	-	-	0
<i>R. hipposideros</i>	-	2	-	-	-	-	-	-	-	2
<i>R. mehelyi</i>	-	-	-	-	-	-	-	-	-	0
<i>T. teniotis</i>	-	-	-	-	-	-	-	-	-	0
<i>M. schreibersii</i>	-	-	-	-	-	-	-	-	-	0
<i>E. isabellinus</i>	42	1	3	64	-	-	2	-	1	113
<i>P. kuhlii</i>	298	47	5	212	67	-	32	2	2	665
<i>P. pipistrellus</i>	128	2	-	143	-	-	10	-	-	283
<i>P. gaisleri</i>	2	-	-	-	-	-	-	-	-	2
<i>M. capaccinii</i>	-	1	-	-	-	-	-	-	-	1
<i>M. punicus</i>	-	7	-	-	-	-	-	-	-	7
Total	470	60	8	419	67	0	44	2	3	

Bat activity was higher at the bridge of Baja Wadi than in the other habitat types in March 2016 ($\chi^2 = 714.23$; d.f. = 2; P < 0.001), August 2016 ($\chi^2 = 625.42$; d.f. = 2; P < 0.001) and February 2017 ($\chi^2 = 70.33$; d.f. = 2; P < 0.001). In the three sessions signals of *P. kuhlii*, *P. pipistrellus* and *E. isabellinus* were recorded, supplemented by *P. gaisleri* in March only.

Bat activity was lower at the artificial basin: 30 contacts / h in March when *P. kuhlii* was again the most active bat followed by *M. punicus*, *E. isabellinus*, *R. hipposideros*, *M. capaccinii* and *P. pipistrellus*, 33 contacts / h in August and 1 contact / h in February of *P. kuhlii* only. However, during this last session the SM2+ Bat detector, recording overnight, detected calls of five species: *R. ferrumequinum*, *R. hipposideros*, *P. kuhlii*, *P. pipistrellus* and *M. capaccinii*.

Bat activity was very low in crop fields, 4 contacts / h of *P. kuhlii* and *E. isabellinus* in March, null in summer when they were harvested, 1.5 contacts / h of *P. kuhlii* and *E. isabellinus* in February.

Discussion

We report for the first time the presence of 11 bat species in Mastouta-Bishshouk region. Nine species were reported from roosts and seven from foraging areas. This supports the need to combine different sampling methods for inventory of bat communities (Flaquer et al., 2007). This bat richness is similar to the previous survey in Jebel Mgilla National Park (Dalhoumi et al., 2014), slightly lower than the more extensive monthly monitoring in Bou Hedma National Park (Dalhoumi et al., 2016), and clearly lower than the 19 bat species reported in Tunisia (Dalhoumi et al., 2011). Except *Plecotus gaisleri*, all the recorded species were reported in this northern part of Tunisia during the last decade (Puechmaille et al., 2012; Zava & Masseti 2007). As all the species of the genus *Plecotus* emit echolocation calls with low intensity they are hardly recorded by bat detectors (e.g. Barataud, 2012),

In roost surveys acoustic recording performed better than captures (7 species vs 5 species). These results confirm that combined methods are also required for bat inventories in roosts (Rnjak et al., 2017). However capture is necessary for a morphological identification when *Rhinolophus euryale* and/or *R. mehelyi* are present (Puechmaille et al., 2012) as the call frequencies of these two species overlap (Weid & von Helversen, 1987; Heller & von Helversen, 1989; Papadatou et al., 2008).

The train abandoned tunnel was the most important roost site found in the study area that lacks natural underground sites. The ceiling of this tunnel hosted a mixed maternity colony of *R. ferrumequinum*, *M. schreibersii* and *M. capaccinii*, confirming the breeding status of the later species in northern Tunisia (Dalhoumi et al., 2011). This tunnel is a roost preserved from human disturbance when the tunnel is flooded. On the contrary, the tunnel of water channel, which is frequently visited by local people and where we only found a little amount of faecal pellets, seems to be a temporary roost.

When foraging, bats were mainly recorded near water, crop fields being deserted after harvesting. High levels of bat activity at the bridge and the basin support previous studies that found higher foraging activity of bats over water than in adjacent habitats (Dalhoumi et al., 2014, 2016; Fukui et al., 2006; Hagen & Sabo, 2011; Grindal et al., 1999; Russo & Jones, 2003; Seibold et al., 2013). Water bodies are known to usually provide a higher density of insect prey than other habitat types (Fukui et al., 2006; Hagen & Sabo, 2011).

Pipistrellus kuhlii, *P. pipistrellus* and *Eptesicus isabellinus* were the most recorded species both in the three sampled sites and in the three surveys. *E. isabellinus* and *P. kuhlii* are the most widespread and the more common bat species in Tunisia (Dalhoumi et al., 2011; Puechmaille et al., 2012) and in the whole Northwest Africa (Aulagnier, 2013; Van

Cakenberghe & Benda, 2013). They were recorded in all sites in Jebel Mghilla National Park (Dalhoumi et al., 2014) and all sampled habitats in Bou Hedma National Park (Dalhoumi et al., 2017 a, b).

We recorded the highest level of activity in late March. The activity was still high in late August, a result contrasting with data from arid central Tunisia (Bou Hedma National Park) where bat activity declined during late summer (Dalhoumi et al., 2016). In late February, bat activity was very low, suggesting that most bats still hibernate according to previous records in caves of Djebel Zaghouan (Deleuil & Labbe, 1955). Such low winter activity was previously reported from northern Algeria (Gaisler, 1984) and Morocco (Aulagnier et al., 2017).

Our survey was conducted only at two seasons and during a few days, so this bat assemblage could be larger if we consider the more recent checklists for northern Tunisia (Dalhoumi et al., 2011; Puechmaille et al., 2012). Three species could also be present in Mastouta-Bishshouk region: *Rhinolophus blasii*, *R. euryale* and *Myotis emarginatus* but these species were quite rarely reported in the country. Restricted so far to northern Tunisia and only reported in four localities (Puechmaille et al., 2012) due to the difficulty to locate roosts, *Hypsugo savii* has been detected at the dam and the mine of Beni Metir (Puechmaille et al., 2012), 40 km far from the study area. Its presence should be further investigated using passive ultrasound detectors near cliffs which offer potential roost to this species.

Apart from the habitat-related activity of bats in this part of Tunisia the main result of this survey is the discovery of a mixed maternity colony, including *M. capaccinii*, in the train abandoned tunnel. This site should require conservation attention as underground roosts of this size are rare in the region and most often suffer human disturbance. Obviously the flooding of the tunnel, except during exceptional drought, offers a natural protection to this roost. The water quality of Majerda Wadi should be improved as this river collect pesticides used in crop fields, factory and domestic sewerages.

The authors thank the Exploralis association and its members who facilitated this study in the mouth of the Sidi Salem Dam and the neighboring regions. This study was funded by MAVA's foundation and the French Funds for the Global Environment. We also thank the General Directorate of Forestry, the General Directorate of Waters (Ministry of Agriculture) and the CRDA of Béja for the licenses provided. We are grateful to reviewers for important comments on the manuscript.

References

- Altringham, J. D. 2011. *Bats. From evolution to conservation*. Second edition. Oxford University Press, Oxford (UK), 1–324.
- Aulagnier, S. 2013. *Eptesicus serotinus*. Common serotine. In: Happold, M., Happold, D. C. D., eds. *Mammals of Africa. Vol. IV. Hedgehogs, shrews and bats*. Bloomsbury Publishing, London (UK), 558–559.
- Aulagnier, S., Haffner, P., Mitchell-Jones, T., Moutou, F., Zima, J. 2009. *Mammals of Europe, North Africa and the Middle East*. A & C Black, London (UK), 1–272.
- Aulagnier, S., Cuzin, F., Thévenot, M. 2017. Chiroptera. In: Aulagnier, S., Cuzin F., Thévenot, M., eds. *Mammifères sauvages du Maroc. Peuplement, répartition, écologie*. Société Française pour l'Etude et la Protection des Mammifères, Paris (France), 117–154.
- Barataud, M. 2012. *Ecologie acoustique des Chiroptères d'Europe. Identification des espèces, études de leurs habitats et comportements de chasse*. Biotope. Muséum National d'Histoire Naturelle. Mèze-Paris (France), 1–344.
- Blanc, M. 1935. *Faune tunisienne. I. Mammifères*. Document dactylographié, Tunis (Tunisia), 1–43.
- Bright, P. 1993. Habitat fragmentation and Prediction for British Mammals. *Mammal Review*, **23** (3–4), 101–111.
- Castro, I. J., Michalski, F. 2014. Effects of logging on bats in tropical forests. *Natureza & Conservação*, **12** (2), 99–105.
- Chriha, S., Sghari, A. 2013. Les incendies de forêt en Tunisie, Séquelles irréversibles de la révolution de 2011. *Méditerranée*, **121**, 87–93.
- Cockrum, E. L. 1976. Order Chiroptera. In: *Mammals of Tunisia*. Unpublished report, 72–126.

- Dalhoumi, R., Aissa P., Aulagnier, S. 2011. Taxonomie et répartition des Chiroptères de Tunisie. *Revue suisse de Zoologie*, **118** (2), 265–292.
- Dalhoumi, R., Hedfi, A., Aissa, P., Aulagnier, S. 2014. Bats of Jebel Mghilla National Park (central Tunisia): first survey and habitat-related activity. *Tropical Zoology*, **27** (2), 53–62.
- Dalhoumi R., Aissa P., Aulagnier S. 2016. Bat species richness and activity in the Bou Hedma National Park (central Tunisia). *Turkish Journal of Zoology*, **40** (6), 864–875.
- Dalhoumi, R., Morellet, N., Aissa, P., Aulagnier, S. 2017 a. Seasonal activity pattern and habitat use by the Isabelline serotine bat (*Eptesicus isabellinus*) in an arid environment of Tunisia. *Acta Chiropterologica*, **19** (1), 141–153.
- Dalhoumi R., Morellet N., Aissa P., Aulagnier S. 2017 b. Seasonal activity pattern and habitat use by the Kuhl's pipistrelle (*Pipistrellus kuhlii*) in an arid environment. *European Journal of Wildlife Research* (in press).
- Deleuil, R., Labbe, A. 1955. Contributions à l'étude des chauves-souris de Tunisie. *Bulletin de la Société de Sciences naturelles de Tunisie*, **8**, 39–55.
- Dietz, C., von Helversen, O., Nill, D. 2007. *Handbuch der Fledermäuse Europas und Nordwestafrikas: Biologie, Kennzeichen, Gefährdung*. Kosmos, Stuttgart (Germany), 1–399.
- Erkert, H. G. 1982. Ecological aspects of bat activity rhythms. In: Kunz, T. H., ed. *Ecology of bats*. Plenum Press, Ney York (NY), 201–242.
- Findley, J. S. 1993. *Bats: a community perspective*. Cambridge University Press, Cambridge (UK), 1–167.
- Flaquer, C. I., Torre, I., Arrizabalaga, A. 2007. Comparison of sampling methods for inventory of bat communities. *Journal of Mammalogy* **88** (4), 526–563.
- Fukui, D., Murakami, M., Nakano, S., Aoi, T. 2006. Effect of emergent aquatic insects on bat foraging in a riparian forest. *Journal of Animal Ecology*, **75** (6), 1252–1258.
- Gadeau de Kerville, H. 1908. *Voyage zoologique en Khroumirie (Tunisie) mai-juin 1906*. Jean-Baptiste Bailliére et Fils, Paris (France), 1–316.
- Gaisler, J. 1984. Bats of Northern Algeria and their winter activity. *Myotis*, **21–22**, 89–95.
- Grindal, S., Morissette, J. L., Brigham, R. M. 1999. Concentration of bat activity in riparian habitats over an elevational gradient. *Canadian Journal of Zoology*, **77** (6), 972–977.
- Hagen, E. M., Sabo, J. L. 2011. A landscape perspective on bat foraging ecology along rivers: does channel confinement and insect availability influence the response of bats to aquatic resources in riverine landscapes? *Oecologia*, **166** (3), 751–760.
- Hartmann, R. 1868. Geographische Verbreitung der im nordöstlichen Afrika wild lebenden Säugetiere. *Zeitschrift der Gesellschaft für Erdkunde zu Berlin*, **3**, 28–69.
- Heller, K. G. & Von Helversen O. 1989. Resource partitioning of sonar frequency bands in rhinolophid bats. *Oecologia* **80** (2), 178–186.
- Hill, J. K., Gray, M. A., Chey, V. K., Benedick, S., Ta watao, N., Hamer, K. C. 2011. Ecological impacts of tropical forest fragmentation: how consistent are patterns in species richness and nestedness? *Philosophical Transactions of the Royal Society B: Biological Sciences*, **366**, 3265–3276.
- Neuweiler, G. 2000. *The biology of bats*. Oxford University Press, New York (NY), 1–310.
- Noblet, J. F., Nefzi, T. 1991. *Les chauves-souris de Tunisie. Etude et protection*. Document dactylographié, 1–5.
- Papadatou, E., Butlin, R. K., Altringham, J. D., 2008. Identification of bat species in Greece from their echolocation calls. *Acta Chiropterologica* **10** (1), 127–143.
- Puechmaille, S. J., Hizem, M. W., Allegrini, B., Abiad, A. 2012. Bat fauna of Tunisia: review of records and new records, morphometrics and echolocation data. *Vespertilio*, **16**, 211–239.
- Racey, P. A. 1998. Ecology of European bats in relation to their conservation. In: Kunz, T. H., Racey, P. A., eds. *Bat biology and conservation*. Smithsonian Institution Press, Washington (DC), 249–260.
- Rebelo, H., Brito, J. C. 2006. Bat guild structure and habitat use in the Sahara desert. *African Journal of Ecology*, **45** (2), 228–230.
- Rnjak, D., Goran, R., Zrncić, V. 2017. Godine bat fauna research in the south-eastern part of Istria (Croatia) in 2013. *Hypsugo*, **2** (1), 1–16.
- Russ, J. 1999. *The bats of Britain and Ireland: echolocation calls, sound analysis, and species identification*. Alana Ecology Books, Belfast (UK), 1–103.
- Russ, J. M., Montgomery, W. I. 2002. Habitat associations of bats in Northern Ireland: implications for conservation. *Biological Conservation*, **108**, 49–58.
- Russo, D., Jones, G. 2002. Identification of twenty-two bat species (Mammalia: Chiroptera) from Italy by analysis of time-expanded recordings of echolocation calls. *Journal of Zoology*, **258** (1), 91–103.
- Russo, D., Jones, G. 2003. Use of foraging habitats by bats in a Mediterranean area determined by acoustic surveys: conservation implications. *Ecography*, **26** (2), 197–209.
- Seibold, S., Buchner, J., Bässler, C., Müller, J. 2013. Ponds in acidic mountains are more important for bats in providing drinking water than insect prey. *Journal of Zoology*, **290** (4), 302–308.
- Van Cakenberghe, V., Benda, P. 2013. *Pipistrellus kuhlii* Kuhl's pipistrelle. In: Happold, M., Happold, D. C.

- D., eds. *Mammals of Africa. Vol. IV. Hedgehogs, shrews and bats.* Bloomsbury Publishing, London (UK), 633–635.
- Weid, R., von Helversen, O. 1987. Ortungsrufe europäischer Fledermäuse beim Jagdflug im Freiland. *Myotis* **25**, 5–27.
- Wickramasinghe, L. P., Harris, S., Jones, G., Vaughan, N. 2003. Bat activity and species richness on organic and conventional farms: impact of agricultural intensification. *Journal of applied Ecology*, **40**, 984–993.
- Zava, B., Masseti, M. 2007. First record of four species of bats from the national park of El Feidja (NW Tunisia) with a note on the non-flying mammals of the Khroumiria region. *Hystrix*, **18** (1) (n. s.), 83–90.

Received 18 October 2017

Accepted 25 October 2018